CLEANSLATE RTU disinfectant cleaner

- Tuberculocidal
- This product kills SARS-CoV-2, which causes COVID-19** on hard non-porous surfaces in just 60 seconds!
- Kills 99.9% Of Germs*. Kills 99.9% of Bacteria*
- Kills the virus that causes the common cold++

ttRhinovirus *when used according to disinfection directions **SARS-Related Coronavirus 2

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. IF SWALLOWED: Call a poison control center or doctor mediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Cleanslate RTU Disinfectant Cleaner Vitex is designed specifically as a general ready-to-use cleaner and disinfectant for use on hard, non-porous surfaces in homes, shower rooms, locker rooms, public restrooms, schools, restaurants, commercial kitchens, transportation terminals, office buildings, retail businesses, and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination. It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets. For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable

ACTIVE INGREDIENTS:

F6591

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	NET CONTENTS: 1 Quart (32 fl. oz.)
dimethyl benzyl ammonium chlorides 0.105%	
n-Alkyl (68% C ₁₀ , 32% C ₁₄)	
dimethyl ethylbenzyl ammonium chlorides 0.105%	
INERT INGREDIENTS <u>99.790%</u>	
TOTAL100.00%	
EPA REG. NO. 1839-83-84179 EPA EST. NO. 84179-NJ-1	4 5 9 1
Sold by: Cleanslate Group, LLC. 1420 East Linden Ave., Linden, NJ 07036	1.5.1

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. DISINFECTION, DEODORIZING AND CLEANING: Remove visible soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens; allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 1 minute. No scrubbing is necessary. Wipe off with a clean cloth, mop, paper towel or sponge or allow to air dry. The product will not leave grit or soap scum. MILDEWSTAT - To control mold and mildew on pre-cleaned, hard, non-porous surfaces spray surface to be treated making sure to wet completely. Let air dry. Repeat application at weekly intervals or when mildew growth appears. FUNGICIDAL ACTIVITY - This product is fungicidal against the pathogenic fungus, Trichophyton interdigitale, formerly Trichophyton mentagrophytes, when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, showe stalls, locker rooms, or other areas commonly contacted by bare feet. **†VIRUCIDAL ACTIVITY** - This product kills on hard, non-porous inanimate surfaces when allowed to remain wet for a 10 minute contact time against; Canine Paryovirus (ATCC VR-2017), Hepatitis A Virus (HAV) (University of Ottawa) and Poliovirus Type 1 (ATCC VR-1000). This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: Human Coronavirus (ATCC VR-740), SARS associated Coronavirus (ZeptoMetrix), Avian Influenza A strain H3N2 (ATCC VR-2072) and Avian Influenza A strain H9N2 (Charles River Laboratories) for 2 minutes. Paramyxovirus (Mumos) (ATCC VR-1438). Rhinovirus type 39 (ATCC VR-340) and Rotavirus (University of Ottawa) for 3 minutes: HIV-1 (associated with AIDS) (Advanced Biotechnologies, Inc.), SARS-Related Coronavirus 2 [SARS-CoV-2] [USA-WA1/2020] [BEI NR-52281] for 1 minute; Hepatitis B Virus (HBV) (Hepadnavirus Testing, Inc.), Hepatitis C Virus (HCV) and Bovine Viral Diarrhea Virus (BVDV) (NVSL) when the treated surface is allowed to remain wet for 1 minute; and Norovirus (Norwalk virus), Feline Calicivirus (University of Ottawa) and Rabies Virus (CDC) for 30 seconds. KILLS HIV-1, HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV), and Hepatitis C Virus (HCV). SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. PERSONAL PROTECTION: When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, and eve coverings, CLEANING PROCEDURES; Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. CONTACT TIME: Allow surface to remain wet for HBV, HCV and HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria. DISPOSAL OF INFECTIOUS MATERIALS: Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal. One-step* cleaning and disinfection of toilet bowls and urinals: Remove water from bowl by forcing water over trap with a swab mop applicator. Press swab mop against side of the bowl to remove excess water from applicator. Apply 1 to 2 ounces of this product evenly, allowing sufficient time for product to cover entire surface of bowl for maximum cleaning. Swab bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1, HBV and HCV 1 minute and flush. If water cannot be removed from toilet, use product on the external surface and above waterline only. To disinfect food processing premises: Before using product food products packaging materials must be removed or carefully protected. For floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1, HBV and HCV 1 minute. Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Visibly soiled areas must be pre-cleaned before using this product. To disinfect food service establishment food contact surfaces: Before using this product food products and packaging materials must be removed or carefully protected. For countertops, appliances, tables, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1, HBV and HCV 1 minute. Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Visibly soiled areas must be pre-cleaned before using this product. Do not use on eating utensils, glassware and cookware.

PRECAUTIONARY STATEMENTS Hazards To Humans And Domestic Animals

CAUTION. KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. PESTICIDE STORAGE: Store in a dry place no lower in temperature than 50°F or higher than 120°F. PESTICIDE DISPOSAL. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or EnvironManage and the hazardous. Marging to the second structure and recap. Agitate vigorously. Follow Pesticide Disposal instructions for rinsate lisposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill. Toll-Free 1888-726-8323 Offices 1908-912-2500 Website I cleanneeds.com + greenexclean.com